

Key Findings (2009):

- In all LHINs, there is no significant difference in attachment rates for those who rate their health as excellent, very good, or good, and those who rate their health as fair or poor.

Attachment by SRH, LHIN

Attachment rates for adults, 16 years of age and older, by self-reported health status and LHIN, 2008-2009									
<i>EXAMPLE INTERPRETATION: IN 2009, 93.2% OF ERIE ST. CLAIR LHIN RESIDENTS WHO REPORTED THEIR HEALTH TO BE EXCELLENT, VERY GOOD AND GOOD WERE ATTACHED TO FAMILY DOCTORS.</i>									
LHIN Number	LHIN Name	Characteristic	Categories	2008			2009		
				Proportion	LCI	UCI	Proportion	LCI	UCI
1	Erie St. Clair	Self-Reported Health Status	Excellent, very good, and good self-reported health	90.2%	87.2%	93.2%	93.2%	90.2%	96.2%
			Fair and poor self-reported health	88.7%	80.7%	96.7%	92.2%	84.9%	99.4%
2	South West	Self-Reported Health Status	Excellent, very good, and good self-reported health	92.9%	90.4%	95.3%	91.6%	88.5%	94.7%
			Fair and poor self-reported health	91.0%	85.5%	96.6%	85.8%	73.0%	98.6%
3	Waterloo Wellington	Self-Reported Health Status	Excellent, very good, and good self-reported health	94.7%	92.7%	96.7%	95.6%	93.7%	97.5%
			Fair and poor self-reported health	94.1%	89.2%	99.0%	92.2%	85.5%	99.0%
4	Hamilton Niagara Haldimand Brant	Self-Reported Health Status	Excellent, very good, and good self-reported health	96.1%	94.3%	97.9%	97.1%	95.5%	98.6%
			Fair and poor self-reported health	98.1%	96.1%	100.0%	97.8%	94.4%	100%
5	Central West	Self-Reported Health Status	Excellent, very good, and good self-reported health	92.4%	89.3%	95.4%	93.4%	90.8%	96.1%
			Fair and poor self-reported health	97.8%	94.9%	100%	95.8%	92.2%	99.5%
6	Mississauga Halton	Self-Reported Health Status	Excellent, very good, and good self-reported health	94.1%	91.6%	96.7%	93.3%	90.3%	96.2%
			Fair and poor self-reported health	98.4%	96.2%	100%	88.6%	79.5%	97.6%
7	Toronto Central	Self-Reported Health Status	Excellent, very good, and good self-reported health	91.4%	88.7%	94.1%	91.5%	88.8%	94.3%
			Fair and poor self-reported health	91.1%	82.9%	99.3%	85.7%	75.0%	96.3%
8	Central	Self-Reported Health Status	Excellent, very good, and good self-reported health	94.6%	92.3%	96.8%	94.5%	92.3%	96.7%
			Fair and poor self-reported health	95.8%	90.8%	100%	100%	Not able to calculate*	
9	Central East	Self-Reported Health Status	Excellent, very good, and good self-reported health	92.6%	89.6%	95.7%	93.5%	91.0%	96.0%
			Fair and poor self-reported health	94.7%	90.5%	99.0%	96.0%	91.3%	100%
10	South East	Self-Reported Health Status	Excellent, very good, and good self-reported health	94.0%	91.7%	96.2%	97.3%	95.5%	99.1%
			Fair and poor self-reported health	89.2%	80.1%	98.3%	96.3%	92.2%	100%
11	Champlain	Self-Reported Health Status	Excellent, very good, and good self-reported health	92.1%	89.5%	94.7%	89.8%	86.3%	93.4%
			Fair and poor self-reported health	90.1%	83.5%	96.7%	90.3%	84.4%	96.2%

Key Findings (2009):

- In all LHINs, there is no significant difference in attachment rates for those who rate their health as excellent, very good, or good, and those who rate their health as fair or poor.

Attachment by SRH, LHIN

Attachment rates for adults, 16 years of age and older, by self-reported health status and LHIN, 2008-2009									
<i>EXAMPLE INTERPRETATION: IN 2009, 93.2% OF ERIE ST. CLAIR LHIN RESIDENTS WHO REPORTED THEIR HEALTH TO BE EXCELLENT, VERY GOOD AND GOOD WERE ATTACHED TO FAMILY DOCTORS.</i>									
LHIN Number	LHIN Name	Characteristic	Categories	2008			2009		
				Proportion	LCI	UCI	Proportion	LCI	UCI
12	North Simcoe Muskoka	Self-Reported Health Status	Excellent, very good, and good self-reported health	91.1%	88.1%	94.1%	94.2%	91.9%	96.4%
			Fair and poor self-reported health	91.2%	84.7%	97.6%	95.2%	90.6%	99.9%
13	North East	Self-Reported Health Status	Excellent, very good, and good self-reported health	87.0%	83.4%	90.6%	83.5%	78.9%	88.1%
			Fair and poor self-reported health	83.9%	76.8%	91.1%	82.3%	73.3%	91.4%
14	North West	Self-Reported Health Status	Excellent, very good, and good self-reported health	86.3%	82.6%	89.9%	87.0%	83.6%	90.4%
			Fair and poor self-reported health	89.0%	82.1%	95.9%	85.6%	77.0%	94.3%
ONTARIO		Self-Reported Health Status	Excellent, very good, and good self-reported health	92.8%	92.1%	93.6%	93.0%	92.3%	93.8%
			Fair and poor self-reported health	93.5%	92.0%	95.0%	92.9%	91.0%	94.8%

Key Findings (2009):

- In the Erie St. Clair, Mississauga Halton and Central LHINs, individuals with one or more chronic conditions are significantly more likely to be attached than those with no chronic conditions.

Attachment by chronic, LHIN

Attachment rates for adults, 16 years of age and older, by chronic conditions and LHIN, 2008-2009									
<i>EXAMPLE INTERPRETATION: IN 2009, 96.8% OF ERIE ST. CLAIR LHIN RESIDENTS WHO HAVE ONE OR MORE CHRONIC CONDITIONS WERE ATTACHED TO FAMILY DOCTORS.</i>									
LHIN Number	LHIN Name	Characteristic	Categories	2008			2009		
				Proportion	LCI	UCI	Proportion	LCI	UCI
1	Erie St. Clair	Select Chronic Condition (1 or more)	No	85.4%	80.7%	90.1%	89.2%	84.4%	94.1%
			Yes	94.4%	91.3%	97.6%	96.8%	94.4%	99.3%
2	South West	Select Chronic Condition (1 or more)	No	93.0%	90.0%	96.1%	87.6%	82.7%	92.4%
			Yes	92.2%	88.9%	95.5%	94.8%	92.3%	97.4%
3	Waterloo Wellington	Select Chronic Condition (1 or more)	No	93.9%	91.2%	96.6%	95.3%	93.0%	97.7%
			Yes	95.6%	93.2%	98.1%	95.1%	92.1%	98.2%
4	Hamilton Niagara Haldimand Brant	Select Chronic Condition (1 or more)	No	96.7%	94.8%	98.5%	95.9%	93.6%	98.3%
			Yes	96.2%	93.8%	98.7%	98.3%	96.9%	99.8%
5	Central West	Select Chronic Condition (1 or more)	No	92.4%	88.7%	96.0%	93.2%	89.8%	96.6%
			Yes	94.7%	91.2%	98.2%	94.9%	92.5%	97.4%
6	Mississauga Halton	Select Chronic Condition (1 or more)	No	92.9%	89.6%	96.3%	89.5%	85.1%	93.9%
			Yes	97.4%	94.8%	99.9%	97.3%	94.8%	99.8%
7	Toronto Central	Select Chronic Condition (1 or more)	No	90.9%	87.6%	94.2%	88.9%	85.1%	92.7%
			Yes	92.5%	88.8%	96.2%	94.0%	90.8%	97.3%
8	Central	Select Chronic Condition (1 or more)	No	93.4%	90.3%	96.4%	92.6%	89.4%	95.9%
			Yes	96.8%	94.5%	99.2%	98.4%	96.9%	99.9%
9	Central East	Select Chronic Condition (1 or more)	No	93.4%	89.8%	96.9%	91.7%	88.3%	95.1%
			Yes	92.5%	88.8%	96.3%	96.6%	93.8%	99.4%

Key Findings (2009):

- In the Erie St. Clair, Mississauga Halton and Central LHINs, individuals with one or more chronic conditions are significantly more likely to be attached than those with no chronic conditions.

Attachment by chronic, LHIN

Attachment rates for adults, 16 years of age and older, by chronic conditions and LHIN, 2008-2009									
<i>EXAMPLE INTERPRETATION: IN 2009, 96.8% OF ERIE ST. CLAIR LHIN RESIDENTS WHO HAVE ONE OR MORE CHRONIC CONDITIONS WERE ATTACHED TO FAMILY DOCTORS.</i>									
LHIN Number	LHIN Name	Characteristic	Categories	2008			2009		
				Proportion	LCI	UCI	Proportion	LCI	UCI
10	South East	Select Chronic Condition (1 or more)	No	91.7%	88.1%	95.3%	96.3%	93.5%	99.1%
			Yes	95.1%	92.3%	97.9%	98.0%	96.3%	99.7%
11	Champlain	Select Chronic Condition (1 or more)	No	90.5%	86.8%	94.3%	89.7%	85.2%	94.2%
			Yes	93.4%	90.5%	96.3%	90.1%	85.8%	94.5%
12	North Simcoe Muskoka	Select Chronic Condition (1 or more)	No	89.5%	85.5%	93.6%	92.8%	89.5%	96.1%
			Yes	93.2%	90.0%	96.4%	96.0%	93.7%	98.2%
13	North East	Select Chronic Condition (1 or more)	No	87.9%	83.3%	92.4%	79.3%	72.8%	85.9%
			Yes	84.7%	80.0%	89.4%	87.4%	82.7%	92.1%
14	North West	Select Chronic Condition (1 or more)	No	84.5%	79.4%	89.6%	82.7%	77.8%	87.6%
			Yes	89.2%	85.6%	92.8%	91.4%	87.6%	95.2%
ONTARIO		Select Chronic Condition (1 or more)	No	92.2%	91.2%	93.2%	91.0%	89.9%	92.1%
			Yes	93.9%	93.0%	94.8%	95.4%	94.6%	96.2%

Key Findings (2009):

- For overall care, compared to Ontario overall, utilization is significantly higher in the Toronto Central and Central LHINs.
- For routine care, compared to Ontario overall, utilization is significantly lower in the HNHB LHIN, and is significantly higher in the Toronto Central LHIN.
- For immediate care, compared to Ontario overall, utilization is significantly lower in the South West and North East LHINs.

Primary Care by LHIN

Proportion of attached adults, 16 years of age and older, who used types of primary care in the last 12 months, by LHIN, 2008-2009								
<i>EXAMPLE INTERPRETATION: IN 2009, 80.2% OF ATTACHED ADULTS IN THE ERIE ST. CLAIR LHIN SAW DOCTORS FOR OVERALL CARE.</i>								
LHIN Number	LHIN Name	Type of utilization	2008			2009		
			Proportion	LCI	UCI	Proportion	LCI	UCI
1	Erie St. Clair	Overall care	82.4%	78.7%	86.2%	80.2%	76.3%	84.1%
		Routine care	70.8%	66.4%	75.2%	70.0%	65.5%	74.6%
		Immediate care	33.5%	29.0%	38.0%	33.8%	29.3%	38.3%
2	South West	Overall care	76.3%	71.6%	80.9%	82.1%	78.3%	85.9%
		Routine care	65.3%	60.4%	70.2%	72.4%	68.1%	76.6%
		Immediate care	26.8%	22.4%	31.2%	29.7%	25.3%	34.1%
3	Waterloo Wellington	Overall care	78.3%	74.4%	82.3%	82.1%	78.1%	86.0%
		Routine care	64.7%	60.5%	69.0%	70.0%	65.7%	74.3%
		Immediate care	32.0%	27.6%	36.3%	33.1%	28.5%	37.7%
4	Hamilton Niagara Haldimand Brant	Overall care	82.8%	79.2%	86.4%	81.2%	77.5%	84.9%
		Routine care	71.2%	67.0%	75.4%	65.5%	61.0%	69.9%
		Immediate care	35.3%	30.8%	39.7%	35.6%	31.4%	39.9%
5	Mississauga Halton	Overall care	83.5%	79.9%	87.2%	85.0%	81.4%	88.6%
		Routine care	69.3%	65.0%	73.6%	70.8%	66.3%	75.2%
		Immediate care	40.7%	36.2%	45.2%	41.6%	36.8%	46.4%
6	Central West	Overall care	86.9%	83.6%	90.1%	85.3%	81.8%	88.8%
		Routine care	76.7%	72.7%	80.7%	74.6%	70.4%	78.8%
		Immediate care	41.0%	36.4%	45.6%	40.4%	35.7%	45.2%
7	Toronto Central	Overall care	86.9%	83.7%	90.1%	88.9%	85.7%	92.0%
		Routine care	75.8%	71.6%	80.0%	78.1%	73.8%	82.4%
		Immediate care	42.8%	38.2%	47.5%	41.5%	36.5%	46.6%
8	Central	Overall care	87.1%	83.9%	90.3%	88.8%	85.6%	91.9%
		Routine care	76.8%	72.8%	80.8%	76.1%	72.0%	80.1%
		Immediate care	39.5%	34.7%	44.3%	38.3%	33.3%	43.2%

Key Findings (2009):

- For overall care, compared to Ontario overall, utilization is significantly higher in the Toronto Central and Central LHINs.
- For routine care, compared to Ontario overall, utilization is significantly lower in the HNH B LHIN, and is significantly higher in the Toronto Central LHIN.
- For immediate care, compared to Ontario overall, utilization is significantly lower in the South West and North East LHINs.

Primary Care by LHIN

Proportion of attached adults, 16 years of age and older, who used types of primary care in the last 12 months, by LHIN, 2008-2009								
<i>EXAMPLE INTERPRETATION: IN 2009, 80.2% OF ATTACHED ADULTS IN THE ERIE ST. CLAIR LHIN SAW DOCTORS FOR OVERALL CARE.</i>								
			2008			2009		
LHIN Number	LHIN Name	Type of utilization	Proportion attached			Proportion attached		
			Proportion	LCI	UCI	Proportion	LCI	UCI
9	Central East	Overall care	83.5%	79.9%	87.2%	81.4%	77.7%	85.2%
		Routine care	68.9%	64.6%	73.3%	69.1%	64.4%	73.7%
		Immediate care	34.2%	29.8%	38.7%	36.5%	31.6%	41.4%
10	South East	Overall care	82.6%	79.1%	86.2%	83.4%	79.7%	87.1%
		Routine care	73.2%	69.2%	77.2%	74.1%	69.9%	78.4%
		Immediate care	31.9%	27.6%	36.2%	30.3%	26.1%	34.5%
11	Champlain	Overall care	86.2%	82.8%	89.7%	86.7%	83.3%	90.2%
		Routine care	77.1%	72.9%	81.2%	75.5%	71.2%	79.8%
		Immediate care	32.0%	27.7%	36.3%	34.2%	29.6%	38.9%
12	North Simcoe Muskoka	Overall care	83.6%	80.1%	87.1%	82.6%	79.3%	85.9%
		Routine care	74.9%	70.8%	79.1%	71.4%	67.0%	75.9%
		Immediate care	29.9%	25.5%	34.3%	30.4%	25.8%	35.1%
13	North East	Overall care	83.8%	80.2%	87.3%	78.8%	74.5%	83.2%
		Routine care	70.5%	66.1%	74.9%	72.0%	67.4%	76.6%
		Immediate care	29.4%	25.1%	33.7%	25.0%	20.9%	29.1%
14	North West	Overall care	78.6%	74.4%	82.8%	80.6%	76.6%	84.7%
		Routine care	70.0%	65.5%	74.6%	71.1%	66.6%	75.6%
		Immediate care	27.6%	23.4%	31.8%	30.3%	25.9%	34.7%
ONTARIO		Overall care	83.8%	82.7%	84.8%	84.1%	83.0%	85.1%
		Routine care	72.3%	71.1%	73.6%	72.3%	71.0%	73.6%
		Immediate care	35.3%	34.0%	36.7%	35.7%	34.3%	37.1%

Key Findings:

- For the attached population, compared to Ontario overall, residents in the South West and North West LHINs are significantly more likely to use emergency rooms, while residents of the Waterloo Wellington LHIN are significantly less likely.
- For the attached population, compared to Ontario overall, residents of the Mississauga Halton and Central East LHINs are significantly more likely to use walk-in clinics while residents of the South West, Waterloo Wellington and South East LHINs are significantly less likely to use walk-in clinics.

Walk-in, ER use by LHIN

Use of walk-in clinics and emergency departments, by attached adults, 16 years of age and older, by LHIN, 2008-2009								
EXAMPLE INTERPRETATION: IN 2009, 21.1% OF ATTACHED ADULTS IN THE ERIE ST. CLAIR LHIN WENT TO EMERGENCY DEPARTMENTS AND 28.8% WENT TO WALK-IN CLINICS.								
			2008			2009		
LHIN Number	LHIN Name	Utilization of specialty services	Proportion attached			Proportion attached		
			Proportion	LCI	UCI	Proportion	LCI	UCI
1	Erie St. Clair	Used emergency departments	26.5%	22.1%	30.9%	21.1%	17.0%	25.1%
		Used walk-in clinics	29.0%	24.5%	33.5%	28.8%	24.5%	33.0%
2	South West	Used emergency departments	25.4%	20.6%	30.2%	25.9%	21.9%	29.9%
		Used walk-in clinics	14.0%	10.7%	17.3%	12.4%	9.1%	15.6%
3	Waterloo Wellington	Used emergency departments	20.1%	16.3%	23.8%	15.8%	12.6%	18.9%
		Used walk-in clinics	13.1%	9.9%	16.2%	12.7%	9.4%	15.9%
4	Hamilton Niagara Haldimand Brant	Used emergency departments	18.2%	14.8%	21.7%	19.1%	15.3%	22.9%
		Used walk-in clinics	20.4%	16.7%	24.0%	24.6%	20.5%	28.8%
5	Central West	Used emergency departments	17.7%	14.1%	21.3%	17.1%	13.4%	20.7%
		Used walk-in clinics	31.0%	26.5%	35.4%	26.3%	22.2%	30.4%
6	Mississauga Halton	Used emergency departments	18.7%	15.0%	22.4%	16.5%	12.9%	20.1%
		Used walk-in clinics	34.1%	29.7%	38.5%	33.7%	29.2%	38.2%
7	Toronto Central	Used emergency departments	16.9%	13.5%	20.3%	20.8%	16.7%	25.0%
		Used walk-in clinics	28.2%	23.7%	32.7%	26.9%	22.3%	31.5%
8	Central	Used emergency departments	17.2%	13.8%	20.6%	16.9%	13.3%	20.5%

Key Findings:

- For the attached population, compared to Ontario overall, residents in the South West and North West LHINs are significantly more likely to use emergency rooms, while residents of the Waterloo Wellington LHIN are significantly less likely.
- For the attached population, compared to Ontario overall, residents of the Mississauga Halton and Central East LHINs are significantly more likely to use walk-in clinics while residents of the South West, Waterloo Wellington and South East LHINs are significantly less likely to use walk-in clinics.

Walk-in, ER use by LHIN

Use of walk-in clinics and emergency departments, by attached adults, 16 years of age and older, by LHIN, 2008-2009								
<i>EXAMPLE INTERPRETATION: IN 2009, 21.1% OF ATTACHED ADULTS IN THE ERIE ST. CLAIR LHIN WENT TO EMERGENCY DEPARTMENTS AND 28.8% WENT TO WALK-IN CLINICS.</i>								
			2008			2009		
LHIN Number	LHIN Name	Utilization of specialty services	Proportion attached			Proportion attached		
			Proportion	LCI	UCI	Proportion	LCI	UCI
	Central	<i>Used walk-in clinics</i>	29.0%	24.6%	33.3%	28.5%	24.0%	33.0%
9	Central East	<i>Used emergency departments</i>	19.8%	16.1%	23.4%	19.7%	15.5%	23.9%
		<i>Used walk-in clinics</i>	30.0%	25.6%	34.3%	33.1%	28.3%	37.9%
10	South East	<i>Used emergency departments</i>	26.3%	22.2%	30.4%	23.2%	19.0%	27.4%
		<i>Used walk-in clinics</i>	13.4%	10.1%	16.7%	12.5%	9.1%	15.9%
11	Champlain	<i>Used emergency departments</i>	24.7%	20.5%	28.9%	25.0%	20.6%	29.5%
		<i>Used walk-in clinics</i>	21.3%	17.2%	25.3%	23.3%	18.7%	28.0%
12	North Simcoe Muskoka	<i>Used emergency departments</i>	25.2%	20.9%	29.5%	22.3%	18.0%	26.5%
		<i>Used walk-in clinics</i>	24.9%	20.7%	29.2%	25.2%	21.0%	29.4%
13	North East	<i>Used emergency departments</i>	28.3%	23.9%	32.7%	25.4%	21.3%	29.5%
		<i>Used walk-in clinics</i>	23.1%	19.1%	27.0%	25.9%	21.2%	30.6%
14	North West	<i>Used emergency departments</i>	25.5%	21.4%	29.6%	30.2%	25.7%	34.7%
		<i>Used walk-in clinics</i>	20.7%	16.5%	24.9%	21.4%	17.2%	25.6%
	ONTARIO	<i>Used emergency departments</i>	20.9%	19.8%	22.0%	20.4%	19.2%	21.5%
		<i>Used walk-in clinics</i>	24.8%	23.6%	26.0%	25.3%	24.0%	26.6%

Key Findings (2009):

- For the attached population, compared to Ontario overall, residents in the Erie St. Clair, North Simcoe Muskoka and North West LHINs are significantly less likely to receive care within one day for immediate care.

Time to utilization by LHIN

Time to see doctor for immediate care, by attached adults, 16 years of age and older, by LHIN, 2008-2009								
<i>EXAMPLE INTERPRETATION: IN 2009, OF THE ATTACHED ADULTS WHO SAW DOCTORS FOR IMMEDIATE CARE, 35.1% SAW THE DOCTOR WITHIN ONE DAY.</i>								
LHIN Number	LHIN Name	Time to utilization	2008			2009		
			Proportion	LCI	UCI	Proportion	LCI	UCI
1	Erie St. Clair	<i>Within 1 day</i>	38.8%	31.4%	46.1%	35.1%	26.8%	43.4%
		<i>2 or more days</i>	61.2%	53.9%	68.6%	64.9%	56.6%	73.2%
2	South West	<i>Within 1 day</i>	43.3%	34.4%	52.3%	47.1%	38.7%	55.4%
		<i>2 or more days</i>	56.7%	47.7%	65.6%	52.9%	44.6%	61.3%
3	Waterloo Wellington	<i>Within 1 day</i>	43.8%	36.0%	51.6%	48.6%	40.1%	57.1%
		<i>2 or more days</i>	56.2%	48.4%	64.0%	51.4%	42.9%	59.9%
4	Hamilton Niagara Haldimand Brant	<i>Within 1 day</i>	40.7%	33.8%	47.5%	48.0%	40.0%	56.0%
		<i>2 or more days</i>	59.3%	52.5%	66.2%	52.0%	44.0%	60.0%
5	Central West	<i>Within 1 day</i>	52.7%	45.0%	60.5%	48.3%	41.1%	55.6%
		<i>2 or more days</i>	47.3%	39.5%	55.0%	51.7%	44.4%	58.9%
6	Mississauga Halton	<i>Within 1 day</i>	55.7%	48.4%	62.9%	51.9%	44.3%	59.5%
		<i>2 or more days</i>	44.3%	37.1%	51.6%	48.1%	40.5%	55.7%
7	Toronto Central	<i>Within 1 day</i>	49.1%	41.3%	56.9%	51.3%	43.8%	58.8%
		<i>2 or more days</i>	50.9%	43.1%	58.7%	48.7%	41.2%	56.2%
8	Central	<i>Within 1 day</i>	49.8%	42.3%	57.4%	49.3%	41.2%	57.5%
		<i>2 or more days</i>	50.2%	42.6%	57.7%	50.7%	42.5%	58.8%
9	Central East	<i>Within 1 day</i>	43.1%	34.7%	51.5%	46.8%	38.3%	55.2%
		<i>2 or more days</i>	56.9%	48.5%	65.3%	53.2%	44.8%	61.7%
10	South East	<i>Within 1 day</i>	34.3%	26.1%	42.6%	48.8%	39.9%	57.7%
		<i>2 or more days</i>	65.7%	57.4%	73.9%	51.2%	42.3%	60.1%

Key Findings (2009):

- For the attached population, compared to Ontario overall, residents in the Erie St. Clair, North Simcoe Muskoka and North West LHINs are significantly less likely to receive care within one day for immediate care.

Time to utilization by LHIN

Time to see doctor for immediate care, by attached adults, 16 years of age and older, by LHIN, 2008-2009								
<i>EXAMPLE INTERPRETATION: IN 2009, OF THE ATTACHED ADULTS WHO SAW DOCTORS FOR IMMEDIATE CARE, 35.1% SAW THE DOCTOR WITHIN ONE DAY.</i>								
LHIN Number	LHIN Name	Time to utilization	2008			2009		
			Proportion	LCI	UCI	Proportion	LCI	UCI
11	Champlain	<i>Within 1 day</i>	39.9%	31.3%	48.6%	44.7%	36.0%	53.3%
		<i>2 or more days</i>	60.1%	51.4%	68.7%	55.3%	46.7%	64.0%
12	North Simcoe Muskoka	<i>Within 1 day</i>	30.2%	22.1%	38.4%	31.5%	23.1%	40.0%
		<i>2 or more days</i>	69.8%	61.6%	77.9%	68.5%	60.0%	76.9%
13	North East	<i>Within 1 day</i>	39.9%	30.8%	49.0%	46.3%	37.8%	54.8%
		<i>2 or more days</i>	60.1%	51.0%	69.2%	53.7%	45.2%	62.2%
14	North West	<i>Within 1 day</i>	33.7%	24.3%	43.0%	30.4%	22.5%	38.3%
		<i>2 or more days</i>	66.3%	57.0%	75.7%	69.6%	61.7%	77.5%
ONTARIO		<i>Within 1 day</i>	44.4%	42.1%	46.8%	46.7%	44.3%	49.1%
		<i>2 or more days</i>	55.6%	53.2%	57.9%	53.3%	50.9%	55.7%